

# Review for the mid-term exam

Math 325  
Spring, 2006

**Part I.** All HW problems of Chapters 1 - 3.

**Part II.** True or False (Some of these problems require calculations)

1. A system of 4 linear equations in three unknowns is always inconsistent.
2. If the 4x4 matrix  $A$  has rank 4, the any linear system with coefficient matrix will have a unique solution.
3. There exists a 5x5 matrix  $A$  of rank 4 such that the system  $A\vec{x} = \vec{0}$  has only the solution  $\vec{x} = \vec{0}$ .
4. Vector  $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  is a linear combination of vectors

$$\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix}$$

5. If  $A$  is a nonzero matrix of the form  $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ , then the rank of  $A$  must be 2.
6. There exists a 2x2 matrix  $A$  such that

$$A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \text{and} \quad A \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

7. If  $\vec{u}$ ,  $\vec{v}$  and  $\vec{w}$  are nonzero vectors in  $R^2$ , then  $\vec{w}$  must be a linear combination of  $\vec{u}$  and  $\vec{v}$ .
8. If  $\vec{u}$  is a linear combination of vectors  $\vec{v}$  and  $\vec{w}$ , and  $\vec{v}$  is a linear combination of vectors  $\vec{p}$ ,  $\vec{q}$ , and  $\vec{r}$ , then  $\vec{u}$  must be a linear combination of  $\vec{p}$ ,  $\vec{q}$ , and  $\vec{r}$  and  $\vec{w}$ .
9. If  $A$  is a 4x3 matrix of rank 3 and  $A\vec{v} = A\vec{w}$  for two vectors  $\vec{v}$  and  $\vec{w}$  in  $R^3$ , then  $\vec{v} = \vec{w}$ .
10. The function  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x - y \\ y - x \end{bmatrix}$  is a linear transformation.
11. Matrix  $\begin{bmatrix} 1/2 & -1/2 \\ 1/2 & 1/2 \end{bmatrix}$  represents a rotation.
12. If  $AB = I_n$  for two  $n \times n$  matrices  $A$  and  $B$ , the  $A$  must be the inverse of  $B$ .
13. The function  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ 1 \end{bmatrix}$  is a linear transformation.
14. Matrix  $\begin{bmatrix} -0.68 & 0.8 \\ -0.8 & -0.6 \end{bmatrix}$  represents a rotation.

15.  $\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}^3 = \begin{bmatrix} 1 & 3k \\ 0 & 1 \end{bmatrix}$  for all real numbers  $k$ .
16. If  $A^2 = I_n$ , then  $A$  (matrix) must be invertible.
17. The equation  $A^{-1} = A$  holds for all 2x2 matrices  $A$  representing a reflection.
18. There exist a 2x3 matrix  $A$  and a 3x2 matrix  $B$  such that  $AB = I_2$ .
19. There exist a 3x2 matrix  $A$  and a 2x3 matrix  $B$  such that  $AB = I_3$ .
20. If the linear system  $A^2\vec{x} = \vec{b}$  is consistent, then the system  $A\vec{x} = \vec{b}$  is consistent.
21. If  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$  are linearly independent vectors in  $R^n$ , then they must form a basis of  $R^n$ .
22. The kernel of any invertible matrix consists of the zero vector only.
23. If  $2\vec{u} + 3\vec{v} + 4\vec{w} = 5\vec{u} + 6\vec{v} + 7\vec{w}$ , then vectors  $\vec{u}, \vec{v}$  and  $\vec{w}$  must be linearly dependent.
24. The column vectors of a 5x4 matrix must be linear dependent.
25. If the kernel of a matrix  $A$  consists of the zero vector only, then the column vectors of  $A$  must be linearly independent.
26. If  $\vec{u}, \vec{v}$  and  $\vec{w}$  are in a subspace  $V$  of  $R^n$ , then the vector  $2\vec{u} + 3\vec{v} + 4\vec{w}$  must be in  $V$  as well.
27. If vectors  $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4$  are linearly independent, then vectors  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  are linearly independent.
28. Matrix  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$  is similar to  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .
29. If a 2x2 matrix  $P$  represents the orthogonal projection onto a line in  $R^2$ , the  $P$  must be similar to the matrix  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .
30. If vectors  $\vec{u}, \vec{v}$  and  $\vec{w}$  are linearly dependent, then vector  $\vec{w}$  must be a linear combination of  $\vec{u},$  and  $\vec{v}$ .
31. If  $A$  and  $B$  are invertible matrices of size  $n \times n$ , then  $AB$  must be similar to  $BA$ .
32. If  $AB = 0$  for two 2x2 matrices  $A$  and  $B$ , then  $BA$  must be the zero matrix as well.
33.  $R^2$  is a subspace  $R^3$ .
34. For every subspace  $V$  of  $R^3$ , there exists a 3x3 matrix  $A$  such that  $V = Im(A)$ .
35. If the kernel of a 5x4 matrix  $A$  consists of the zero vector alone, and if  $AB = AC$  for two 4x5 matrix  $B$  and  $C$ , the matrices  $B$  and  $C$  must be equal.